

# Atmospheric Modeling and Winds

D. M. Kass

J. T. Schofield

(JPL/Caltech)

MER 3<sup>d</sup> Landing Site Workshop

March 26, 2002

# Atmospheric Modeling

## Atmospheric profiles

Temperature, Pressure and Density

Surface (-4.181 to -1.3 km) to 200 km

Monte-Carlo profile families

Moderate dust load ( $\tau \sim 0.3$ )

Based on:

- Ames MGCM results
- TES atmospheric retrievals
- Viking Lander pressure measurements
- Pathfinder surface meteorological measurements

Limits landing elevation to -1.3 km or lower

## Near Surface Atmosphere

Diurnal cycle

winds

Near-surface and surface temperature

Limits the mission lifetime

Limits the science return

May limit equipment survivability

Wind Modeling for Entry, Descent, and Landing (EDL)

# Mesoscale Wind Modeling

Almost no wind data for near-surface equatorial Mars

## Mesoscale Models

- Model meteorological phenomena

  - few km to 100 km scale

  - some important phenomena unresolved

- 3-D dynamical atmospheric models

  - track pressure, temperature, wind vectors...

  - non-hydrostatic equations

  - slope illumination and shadowing

- Use nested grids for high resolution

  - cover area of interest

  - necessary context at lower resolution

- Computationally very expensive

# Mesoscale Atmosphere Models

Use 2 models to check validity

MRAMS: Scott Rafkin (SJSU)

Based on terrestrial RAMS (U Colorado)

Uses Ames MGCM as boundary condition

Kinetic energy conserving grid

~ 1.5 km highest horizontal grid spacing

Very high resolution Large Eddy Simulation (LES)

Mars MM5: Anthony Toigo (Cornell) Mark Richardson (Caltech)

Based on terrestrial MM5 (U Penn.)

Uses GFDL MGCM as boundary condition

Vorticity conserving grid

~ 600 m highest horizontal grid spacing

Interactive dust transport

# EDL Wind Analysis

## Qualitative Study

Understand site setting

## Engineering parameterization

- Effective (DC) mean wind field  
Exponential weighting function
- Scaling MPF shear/turbulence model  
MPF model reasonable, but not ideal  
Long wavelength by Fourier analysis  
Short wavelength by scaling TKE

## Engineering wind profiles

Allows use of EDL Monte-Carlo modeling

Random/selected mesoscale profiles

Add high frequency turbulence

# Peer Review

Successful peer review on March 8, 2002

Panel: R. Zurek, J. Barnes, D. Crisp, J. Murphy,  
R. Pielke, N. Renno

Also attended by project management, scientist  
and engineers

## Review Results:

Models are reasonable

Consistent with atmospheric physics principles

Consistent with expectations

season, latitude and topography

Model intercomparison

Agreement with Pathfinder meteorological data

Analysis techniques are appropriate

Site to site differences are significant

# Hematite Overview

Little topography

Low background wind

inbetween the tropical jets

Highly convective

small craters enhance convection


Thick boundary layer ( $\geq 5$  km)

Significant updrafts and downdrafts

|                           | MRAMS      | Mars MM5  |
|---------------------------|------------|-----------|
| Effective (DC) Wind (m/s) | $4 \pm 2$  | $4 \pm 2$ |
| Upward mean wind (m/s)    | 2.5(1.9)   | 1.4       |
| Downward mean wind (m/s)  | -1.1(-1.5) | -1.7      |
| MPF Scale Factors         |            |           |
| Shear                     | 0.4(0.3)   | 0.2       |
| Average Turbulence        | 0.7        |           |
| Peak Turbulence           | 1.2        |           |

- Values in parenthesis are from the LES
- Shear is long wavelength variability.
- Turbulence is short wavelength variability.
- Mean Turbulence is over convective boundary layer

01-125.13:00 sigma=0.8575





## Gusev Overview

In southern (eastward) tropical low-level jet

Strong crater rim upwelling and interior subsidence

modified by global circulation

eastward cross crater confined flow

significant wind shear


Thin convective boundary layer

locally enhanced turbulence

Strong nighttime katabatic flow

|                           | MRAMS     | Mars MM5    |
|---------------------------|-----------|-------------|
| Effective (DC) Wind (m/s) | $7 \pm 2$ | $3 \pm 0.6$ |
| Upward mean wind (m/s)    | 0.4       | 0.3         |
| Downward mean wind (m/s)  | -0.2      | -0.3        |
| MPF Scale Factors         |           |             |
| Shear                     | 0.9       | 0.5         |
| Average Turbulence        | 1.8       |             |
| Peak Turbulence           | 2.1       |             |

- Shear is long wavelength variability.
- Turbulence is short wavelength variability.
- Mean Turbulence is over convective boundary layer



| Guse     |                             | grid 4                        |            |       |       |      |
|----------|-----------------------------|-------------------------------|------------|-------|-------|------|
| y =      | -2. km                      | 295920.00s [0200 UTC +4 sols] | min        | max   | inc   | lab* |
| contours | speed (m/s)                 |                               | 0.7526     | 28.14 | 2.000 | 1e 0 |
| contours | turb kinetic energy (m2/s2) |                               | 0.4995E-03 | 14.35 | 1.000 | 1e 0 |
| vectors  | →                           | 10 m/s horiz 0.21 m/s vert    | 0.7462E-01 | 25.72 |       |      |

# Melas Overview

## Canyon driven circulation

- nighttime down canyon flow

- daytime up-canyon flow

- calm during flow reversal

## Large up-canyon (westerward) flow

- enhanced by westward tropical jet

- enhanced by thermal tide

- enhanced by Tharsis katabatic venting

- models disagree on timing

## Extreme up-wall local flow


- canyon center subsidence

- depressed convective boundary layer

- sideways thermal “venting”

- maintain remnant down-canyon layer

|               |                               |            |            |
|---------------|-------------------------------|------------|------------|
| Melas         |                               | grid 4     |            |
| x = -2243. km | 43164.00s [1910 UTC +0 solis] | min        | max        |
| longitude     | speed (m/s)                   | 0.0000     | 43.00      |
| latitude      | topo (m)                      | 0.0000E+00 | 0.0000E+00 |
| → vectors     | 5 m/s horiz 0.30 m/s vert     | 0.4874E-01 | 44.35      |
|               |                               |            | lab*       |



## Melas Statistics

|                           | MRAMS      | Mars MM5        |
|---------------------------|------------|-----------------|
| Effective (DC) Wind (m/s) | $14 \pm 5$ | $1.3^* \pm 0.7$ |
| Upward mean wind (m/s)    | 0.7        | 0.1             |
| Downward mean wind (m/s)  | -0.8       | -0.1            |
| MPF Scale Factors         |            |                 |
| Shear                     | 0.8        | 0.5             |
| Average Turbulence        | 1.6        |                 |
| Peak Turbulence           | 2.8        |                 |

\* Speeds are significantly higher 2 hours later ( $6 \pm 3$  m/s)

- Shear is long wavelength variability.
- Turbulence is short wavelength variability.
- Mean Turbulence is over convective boundary layer

## Isidis Overview

Mesoscale model runs in progress

Locally flat plain

Likely to have significant convection

Possibly modified by tropical jet

Large topographic relief to south

Relatively close ( $< 50$  km)

May drive katabatic winds

Could generate remnant nighttime downslope flow

May be affected by mid-latitude baroclinic storms

Basin has many local dust storms

Best guess is intermediate between Hematite and Gusev

## Summary

|                   | Hematite   |           | Gusev     |             | Melas      |                 |
|-------------------|------------|-----------|-----------|-------------|------------|-----------------|
|                   | MRAMS      | Mars MM5  | MRAMS     | Mars MM5    | MRAMS      | Mars MM5        |
| Wind Speed (m/s)  |            |           |           |             |            |                 |
| Horizontal        | $4 \pm 2$  | $4 \pm 2$ | $7 \pm 2$ | $3 \pm 0.6$ | $14 \pm 5$ | $1.3^* \pm 0.7$ |
| Upward            | 2.5(1.9)   | 1.4       | 0.4       | 0.3         | 0.7        | 0.1             |
| Downward          | -1.1(-1.5) | -1.7      | -0.2      | -0.3        | -0.8       | -0.1            |
| MPF Scale Factors |            |           |           |             |            |                 |
| Shear             | 0.4(0.3)   | 0.2       | 0.9       | 0.5         | 0.8        | 0.5             |
| Turbulence        |            |           |           |             |            |                 |
| Average           | 0.7        |           | 1.8       |             | 1.6        |                 |
| Peak              | 1.2        |           | 2.1       |             | 2.8        |                 |

\* Speeds are significantly higher 2 hours later ( $6 \pm 3$  m/s)

- Values in parenthesis are from the LES
- Shear is long wavelength variability.
- Turbulence is short wavelength variability.
- Mean Turbulence is over convective boundary layer